

STAT 545: Categorical Data Analysis (Part II)

Liang Li

Department of Biostatistics
The University of Texas MD Anderson Cancer Center

LLi15@mdanderson.org

Fall 2015 at Rice University

Overview of Part II of this class

Oct 19, 2015 to December 2, 2015. There will be homework/projects and a final exam

- Regression model for binary data
- Regression model for ordinal data
- Regression model for counts data
- Extensions of standard regression models for categorical data
- Marginal models for longitudinal categorical data
- Conditional models for longitudinal categorical data

Logistic Regression

Logistic Regression Model

$$\begin{aligned}\pi(x) &= P(Y = 1|X = x) = 1 - P(Y = 0|X = x) \\ &= \text{expit}(\alpha + \beta x) = \frac{\exp(\alpha + \beta x)}{1 + \exp(\alpha + \beta x)} \in (0, 1)\end{aligned}$$

$$\text{logit}[\pi(x)] = \log \frac{\pi(x)}{1 - \pi(x)} = \alpha + \beta x \in (-\infty, \infty)$$

- ① Binary outcome; for binomial outcome, the model is similar
- ② Interpretation of β (log odds ratio)
- ③ Simple visual model checking by grouping (§ 5.1.2)
- ④ Logistic regression with retrospective studies (§ 5.1.4)
- ⑤ Model fitting through maximum likelihood estimation (§ 5.5)
- ⑥ Inference about model parameters and probabilities (§ 5.2.1)
- ⑦ Checking goodness of fit (§ 5.2.5)

The (log) odds ratio and its interpretation

$$\text{logit} [\pi(\mathbf{x})] = \alpha + \beta x$$



$$\text{logit} [\pi(\mathbf{x})] = \alpha + \beta_1 x_1 + \beta_2 x_2 + \dots \beta_p x_p$$

Simple visual model checking by grouping

- 1 Group the (continuous) covariate into 10 categories by cutoffs at the quantiles, with n_i subjects in each group ($i = 1, 2, \dots, 10$)
- 2 Calculate the average covariate within each group (\bar{x}_i)
- 3 Calculate the proportion of $Y = 1$ within each group (\bar{y}_i)
- 4 Plot logit of \bar{y}_i vs. \bar{x}_i . It should be approximately a straight line
- 5 Note: may need correction when $\bar{y}_i = 0$ or 1.

$$\log \frac{\bar{y}_i}{n_i - \bar{y}_i} \Rightarrow \log \frac{\bar{y}_i + 0.5}{n_i - \bar{y}_i + 0.5}$$

- 6 Only work with a single covariate

Logistic regression with retrospective studies (§ 5.1.4)



Model fitting through maximum likelihood estimation (§ 5.5)



Inference on parameters and probabilities (§ 5.2.1)

Test $H_0 : \beta = 0$ in logistic model $\text{logit} [\pi(x)] = \alpha + \beta x$

- ① Wald, Likelihood ratio, and Score tests are applicable (§ 1.3.3)
- ② The predicted probability and **its confidence interval**



Checking goodness of fit (§ 5.2.3)

$$\text{logit} [\pi(\mathbf{x})] = \alpha + \beta_1 x_1 + \beta_2 x_2$$

- 1 Visual checking through grouping (works best with a single covariate)
- 2 Adding interactions, quadratic terms, etc., and testing for significance or looking at AIC/BIC: problematic but widely used
- 3 Making the model more flexible by using splines
- 4 Global goodness of fit checking by *Hosmer & Lemeshow test*

$$\sum_{i=1}^g \frac{\left(\sum_j y_{ij} - \sum_j \hat{\pi}_{ij} \right)^2}{n_i \left(\sum_j \hat{\pi}_{ij} / n_i \right) \left[1 - \left(\sum_j \hat{\pi}_{ij} \right) / n_i \right]} \sim \chi_{g-2}^2$$

- A large value of any global fit statistic merely indicates *some* lack of fit but provides no insight about its nature

Logistic models with categorical predictors (§ 5.3)

- When there is a single categorical predictor, the data can be arranged in an $I \times 2$ contingency table (e.g., Table 5.3)
- When the categories are unordered (e.g., nominal data), the (saturated) model is $\text{logit}(\pi_i) = \beta_i$ ($i = 1, 2, \dots, I$), with I unknown parameters.
- We may write the model as $\text{logit}(\pi_i) = \alpha + \beta_i$ with set-to-zero constraint $\beta_1 = 0$ or sum-to-zero constraint $\sum_i \beta_i = 0$
- The model for subject j ($j = 1, 2, \dots, n$) is $\text{logit}(\pi_j) = \alpha + \sum_{i=1}^I \beta_i 1\{j \in \text{group } i\}$
- When the categories are ordered (e.g., ordinal data), we may assume that $\text{logit}(\pi_i) = \alpha + \beta x_i$
 - The number of parameters reduced with the linear assumption.
 - Be careful about coding x_i ($i=1,2,\dots,I$): (1,2,3) or (1,4,9)?
 - Treat the x_i like a continuous variable.

Cochran-Armitage Trend Test (§ 5.3.5)


- Developed by Armitage (1955) and Cochran (1954) for $I \times 2$ tables with ordered rows
- They used a linear probability model $\pi = \alpha + \beta x_i$
- It is a chi-square test of the independence between rows and columns under the linear assumption. $H_0 : \beta = 0$.
- This test is equivalent to the score statistic for testing $H_0 : \beta = 0$ in the linear logit model.
- Using directed models can improve inferential power
 - If the trend is indeed linear, making use of the linear trend (as in Cochran-Armitage test) is more powerful than not making use of the linear trend (as in $\text{logit}(p_i) = \beta_i$)

Model Selection (§ 6.1)

The data set is $\{Y_i, X_{1i}, X_{2i}, \dots, X_{pi}; i = 1, 2, \dots, n\}$. The logistic regression model is


$$\pi(\mathbf{X}_i) = \text{expit}(\beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_p X_{pi})$$

The p covariates include interactions, quadratic terms, etc. We want to retain only the predictive covariates in the model.

- Model selection is both science and art
- The same principles that you learned in linear model class still apply
- Two goals: (1) complex enough to fit the data well; (2) relatively simple to interpret (avoid overfitting) 
- Confirmatory studies vs. exploratory studies

How many covariates can be included in the model?

$Y_i \sim \text{Bernoulli}$ with $\pi(\mathbf{X}_i) = \text{expit}(\beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_p X_{pi})$

- The effective sample size of a logistic regression is either $\sum_i Y_i$ or $n - \sum_i Y_i$, whichever is smaller
- **The rule of thumb:** no more than the effective sample size divided by 10 (or, 10 events per covariate)
- Including too many covariates may cause non-convergence 
- Avoid multicollinearity, as in linear regression (📖 Page 209, Table 6.1)
 - The overall test is highly significant ($p < 0.0001$)
 - The individual covariates are, in general, not very significant due to the multicollinearity between the horseshoe crab's width and weight ($r = 0.887$)


Forward, backward, and stepwise model selection

$Y_i \sim \text{Bernoulli with } \pi(\mathbf{X}_i) = \text{expit}(\beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_p X_{pi})$

- Forward procedure: (1) start with just the intercept (2) at each step, add the covariate with the smallest p-value in likelihood ratio or Wald test (3) stop when no more significant covariate is available (However, it can stop prematurely due to lack of power)
- Stepwise procedure: at each step, retest the significance of the terms added at previous stages
- Backward procedure: (1) start with full model (2) at each step, remove the covariate with the largest p-value (3) stop when all remaining covariates are significant. (However, full model may not be stable)
- The dummy variables for a single categorical covariate should be added or removed together (likelihood ratio test); do not place an interaction in the model without the main effect terms
- SAS PROC LOGISTIC offers additional entry and exit p-value criteria

Further comment on forward, backward, and stepwise model selection

$Y_i \sim \text{Bernoulli}$ with $\pi(\mathbf{X}_i) = \text{expit}(\beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_p X_{pi})$




-  Page 211, Table 6.2 illustrates
 - three-way interaction is usually not significant (e.g., lack of power) and not desirable (hard to interpret)
 - dropping multiple covariates at once using likelihood ratio test (LRT) or dropping them one at a time (Wald or LRT)
- All these procedures are not rigorously justified (*ad hoc*); use with caution!
- Modern approaches are available (LASSO, bagging, etc.)
- Philosophically, there is no such thing as “the correct model” or “the true model”: ALL MODELS ARE WRONG, SOME ARE USEFUL — George Box

Akaike Information Criterion (AIC)

Select the model with smaller AIC or BIC (L : maximized log likelihood; m : number of parameters in the model; n : sample size)

$$AIC = -2L + 2m$$


$$BIC = -2L + \log(n)m$$

- **Rationale:** Including more covariates will always include the log likelihood, but may cause overfitting; so we put a “penalty” by adjusting for the size of the model. There are mathematical reasons why the penalty must take this form.
- Other penalties are available: HQ, DIC, etc.
- BIC puts more penalty on larger model, and therefore tends to select the simpler model  Page 213
- Like scatter plot smoothing, the “desired” amount of penalty is a somewhat subjective choice 
- Need a comprehensive assessment of AIC/BIC, significance, residuals, scientific rationale, parsimony and interpretability, etc. 

Residuals: Pearson, Deviance, Standardized

Let y_i denote the binomial outcome for n_i trials at setting i of the explanatory variables, $i = 1, 2, \dots, N$. Let $\hat{\pi}_i$ denote the model estimate of $P(Y = 1)$.



- Pearson residual is like the residual for linear regression, but with standardization
- Deviance residual is motivated from the likelihood and deviance (which resembles the sum of squares in linear regression)
- Standardized residual has an approximate $N(0, 1)$ distribution and is the one that we usually use, BUT:
 - use it with grouped data (binomial instead of binary).  Page 217, Table 6.5

Influence diagnosis for logistic regression

- A single observation can have a much more exorbitant influence in linear regression than in logistic regression, since linear regression has no bound on the distance of y_i from the expected value.
- Points that have extreme predictor values need not have high leverage. In fact, the leverage can be relatively small if $\hat{\pi}_i$ is close to 0 or 1.

Predictive power of a logistic regression model: pseudo R^2

- For linear regression $Y_i = \mathbf{X}_i^T \boldsymbol{\beta} + \epsilon_i$, the R^2 is

$$R^2 = 1 - \frac{\sum_i (Y_i - \mathbf{X}_i^T \hat{\boldsymbol{\beta}})^2}{\sum_i (Y_i - \bar{Y})^2}$$

- For logistic regression, the analog

$$1 - \frac{\sum_i (Y_i - \hat{\pi}_i)^2}{\sum_i (Y_i - \bar{Y})^2}$$

may not be nondecreasing as the model gets more complex
(undesirable)




Predictive power of a logistic regression model: pseudo R^2

For logistic regression, a more widely used measure is the pseudo R^2 of McFadden (1974): $\frac{L_M - L_0}{L_S - L_0} = 1 - \frac{L_M}{L_0}$

$$L = \log \prod_{i=1}^N [\pi_i^{y_i} (1 - \hat{\pi}_i)^{1-y_i}] = \sum_{i=1}^N [y_i \log \hat{\pi}_i + (1 - y_i) \log(1 - \hat{\pi}_i)]$$

- L_M is the log likelihood evaluated at the MLE $\hat{\pi}_i = \text{expit}(\mathbf{X}_i^T \hat{\beta})$
- L_0 is the log likelihood evaluated under the MLE of the null model:
 $\hat{\pi}_i = N^{-1} \sum_i y_i$
- L_S is the log likelihood evaluated under the saturated model with
 $\hat{\pi}_i = y_i$. $L_S = 0$

Receiver Operative Characteristics (ROC) curve

- $Y_i = 0$ (non disease) or 1 (disease). Estimated viral load $\hat{\pi}_i \in (0, 1)$. We classify the subject as a case ($Y = 1$) when $\hat{\pi} > c$ and control ($Y = 0$) when $\hat{\pi} \leq c$.
- Sensitivity $P(\hat{\pi} > c | Y = 1) \leftarrow \frac{\sum_i 1\{\hat{\pi}_i > c\}}{\sum_i Y_i}$ 
- Specificity $P(\hat{\pi} \leq c | Y = 0) \leftarrow \frac{\sum_i 1\{\hat{\pi}_i \leq c\}}{\sum_i (1 - Y_i)}$ 
- ROC curve  p225
- The area under the ROC curve (AUC) is reported as c-statistic in SAS PROC LOGISTIC. It is a number between 0 and 1. $AUC = 0.5$ is like flipping a coin. So $AUC < 0.5$ is unlikely. Good classification requires $AUC > 0.80$ (excellent, > 0.9).

ROC Curve




Figure 6.4 ROC curve (from SAS PROC LOGISTIC) for logistic regression model estimating the probability a crab has satellites, using width and color predictors.

Cochran-Mantel-Haenszel Test (§ 6.4)

- Study the association between a treatment variable (e.g., binary) and a disease outcome (e.g., binary) after adjusting for a possibly confounding variable (e.g., categorical or continuous but grouped) that might influence that association
- Example in Table 6.9: multicenter randomized clinical trial comparing treatment vs. placebo on a binary outcome (cured vs. not)
- The logistic regression approach ($i = 1, 2; k = 1, 2, \dots, K; x_i = 1$ or 2):

$$\pi_{ik} = P(Y = 1 | X = i, Z = k)$$

$$\text{logit}(\pi_{ik}) = \alpha + \beta x_i + \beta_k^Z$$

- Test $H_0: \beta = 0$ using Wald or likelihood ratio test
- What if there is interaction between X and Z , i.e., β depends on Z ?

Table 6.9

226

BUILDING, CHECKING, AND APPLYING LOGISTIC REGRESSION MODELS

Table 6.9 Clinical Trial Relating Treatment to Response for Eight Centers, with Expected Value and Variance (of Success Count for Drug) Under Conditional Independence

Center	Treatment	Response		Odds Ratio	μ_{11k}	$\text{var}(n_{11k})$
		Success	Failure			
1	Drug	11	25	1.19	10.36	3.79
	Control	10	27			
2	Drug	16	4	1.82	14.62	2.47
	Control	22	10			
3	Drug	14	5	4.80	10.50	2.41
	Control	7	12			
4	Drug	2	14	2.29	1.45	0.70
	Control	1	16			
5	Drug	6	11	∞	3.52	1.20
	Control	0	12			
6	Drug	1	10	∞	0.52	0.25
	Control	0	10			
7	Drug	1	4	2.0	0.71	0.42
	Control	1	8			
8	Drug	4	2	0.33	4.62	0.62
	Control	6	1			

Source: Beitler and Landis (1985).

often medical centers or clinics; or they may be levels of age or severity of the condition.

Cochran-Mantel-Haenszel Test (§ 6.4)

Data from Center k ($k = 1, 2, \dots, K$)

	cured	not	Total
Treatment	n_{11k}	n_{12k}	n_{1+k}
placebo	n_{21k}	n_{22k}	n_{2+k}
Total	n_{+1k}	n_{+2k}	n_{++k}


- Test H_0 : Treatment and outcome independent conditional on center
- Both the treatment (row) and outcome (column) totals fixed, $n_{11k} \sim$ hypergeometric distribution
- Under the null, the hypergeometric mean and variance of n_{11k} are

$$\mu_{11k} = E(n_{11k}) = n_{1+k}n_{+1k}/n_{++k}$$

$$\text{var}(n_{11k}) = n_{1+k}n_{2+k}n_{+1k}n_{+2k}/[n_{++k}^2(n_{++k} - 1)]$$

- The CMH statistic is $CMH = \frac{[\sum_k (n_{11k} - \mu_{11k})]^2}{\sum_k \text{var}(n_{11k})}$, which has a large sample chi-squared null distribution with $df = 1$.

Cochran-Mantel-Haenszel Test vs. Logistic Regression


- When the sample size per center (also called strata) is moderately large, the two produce similar results (CMH is a score test of the logistic model)
- When the number of strata is large (like matched pairs data), the logistic regression does not apply but CMH still applies
- A point estimator of the overall odds ratio is available from CMH 

$$\hat{\theta}_{CMH} = \frac{\sum_k (n_{11k}n_{22k}/n_{++k})}{\sum_k (n_{12k}n_{21k}/n_{++k})} = \frac{\sum_k n_{++k}p_{11|k}p_{22|k}}{\sum_k n_{++k}p_{12|k}p_{21|k}}$$

- CMH is the standard method for stratified analysis of categorical data, applicable to $I \times J \times K$ contingency table
- If the treatment effect differs across strata (interaction), use logistic model with interaction

Quasi-complete Separation in Logistic Regression

- Be careful about excessively large (or small) odds ratios or excessively large standard errors
 - multi-collinearity; remove one of the correlated covariates
 - complete or quasi-complete separation
- Complete separation: there exists a vector \mathbf{b} such that $\mathbf{b}^T \mathbf{x}_i > 0$ whenever $y_i = 1$ and $\mathbf{b}^T \mathbf{x}_i < 0$ whenever $y_i = 0$. (more likely with continuous covariates)
- Quasi-complete separation: $\mathbf{b}^T \mathbf{x}_i \geq 0$ whenever $y_i = 1$ and $\mathbf{b}^T \mathbf{x}_i \leq 0$ whenever $y_i = 0$. (more likely with categorical covariates)
- Not a problem with linear regression



Regression Models for Multinomial Data

Models for Multinomial Responses (§ 8)

- Nominal data vs. ordinal data: presence or absence of intrinsic order
- For nominal outcome variable $Y = 1, 2, \dots, J$. We need to model $\pi_j(\mathbf{x}) = P(Y = j|\mathbf{x})$ under constraint $\sum_j \pi_j(\mathbf{x}) = 1$.
- Y follows a multinomial distribution with probabilities $\{\pi_1(\mathbf{x}), \dots, \pi_J(\mathbf{x})\}$.
- Baseline-category logit model (e.g., pick J as the baseline/reference category)

$$\log \frac{\pi_j(\mathbf{x})}{\pi_J(\mathbf{x})} = \alpha_j + \beta_j^T \mathbf{x} \quad , \quad j = 1, 2, \dots, J - 1$$


- These $J - 1$ equations determine parameters for logits with other pairs of response categories, as well as the response probabilities:

$$\log \frac{\pi_a(\mathbf{x})}{\pi_b(\mathbf{x})} = \log \frac{\pi_a(\mathbf{x})}{\pi_J(\mathbf{x})} - \log \frac{\pi_b(\mathbf{x})}{\pi_J(\mathbf{x})}$$

Models for Multinomial Responses: baseline-category logit model

- The constraint leads to: $\pi_J(\mathbf{x}) = \frac{1}{1 + \sum_{j=1}^{J-1} \exp(\alpha_j + \beta_j^T \mathbf{x})}$
- The probability for the j -th category ($j = 1, 2, \dots, J-1$):

$$\pi_j(\mathbf{x}) = \frac{\exp(\alpha_j + \beta_j^T \mathbf{x})}{1 + \sum_{j=1}^{J-1} \exp(\alpha_j + \beta_j^T \mathbf{x})}$$

- With more than two response categories ($J > 2$), the probability of a given category need not continuously increase or decrease (e.g., $\pi_j(\mathbf{x})$ may not be a monotone function of \mathbf{x}) 
- The model is fit by maximum likelihood. Let $\mathbf{y}_i = (y_{i1}, y_{i2}, \dots, y_{iJ})$, where $y_{ij} = 1$ when the response is in category j and 0 otherwise, so that $\sum_j y_{ij} = 1$. The log likelihood is:

$$\sum_{i=1}^n \log \left[\prod_{j=1}^J \pi_j(\mathbf{x}_i)^{y_{ij}} \right]$$

Models for Ordinal Responses (§ 8.2)

- $Y = 1, 2, \dots, J$. We need to model $\pi_j(\mathbf{x}) = P(Y = j|\mathbf{x})$ under constraint $\sum_j \pi_j(\mathbf{x}) = 1$.
- Due to the intrinsic ordering of the response categories, we model the cumulative probabilities

$$P(Y \leq j|\mathbf{x}) = \pi_1(\mathbf{x}) + \dots + \pi_j(\mathbf{x})$$

- The cumulative logits are defined as:


$$\text{logit}[P(Y \leq j|\mathbf{x})] = \log \frac{P(Y \leq j|\mathbf{x})}{1 - P(Y \leq j|\mathbf{x})} = \log \frac{\pi_1(\mathbf{x}) + \dots + \pi_j(\mathbf{x})}{\pi_{j+1}(\mathbf{x}) + \dots + \pi_J(\mathbf{x})}$$

- Cumulative logit model

$$\text{logit}[P(Y \leq j|\mathbf{x})] = \alpha_j + \beta^T \mathbf{x} \quad , \quad j = 1, 2, \dots, J - 1$$

- The cumulative logit is monotone in \mathbf{x} ; this feature not available in baseline-category logit model

Proportional Odds Model



- Proportional odds model $\text{logit}[P(Y \leq j|\mathbf{x})] = \alpha_j + \beta^T \mathbf{x}$, where $j = 1, 2, \dots, J - 1$
- Need to assume the same β for each logit. Therefore, *cumulative logit model* is also called the *proportional odds model*

$$\text{logit}[P(Y \leq j|\mathbf{x}_1)] - \text{logit}[P(Y \leq j|\mathbf{x}_2)] = \beta^T (\mathbf{x}_1 - \mathbf{x}_2)$$

- Cumulative odds ratio $\exp(\beta)$ does not depend on j , i.e., $\beta_j \equiv \beta, \forall j$
- We cannot make the model more generalizable by letting replacing β with β_j : the different cumulative probabilities may cross, which is

impossible

Proportional Odds Model




Figure 8.4 Uniform odds ratios AD/BC whenever $x_1 - x_2 = 1$, for all binary collapsings of the response in cumulative logit model of proportional odds form.

likelihood function is

$$\begin{aligned} \prod_{i=1}^n \left[\prod_{j=1}^J \pi_j(\mathbf{x}_i)^{y_{ij}} \right] &= \prod_{i=1}^n \left\{ \prod_{j=1}^J [P(Y \leq j | \mathbf{x}_i) - P(Y \leq j-1 | \mathbf{x}_i)]^{y_{ij}} \right\} \\ &= \prod_{i=1}^n \left\{ \prod_{j=1}^J \left[\frac{\exp(\alpha_j + \beta^T \mathbf{x}_i)}{1 + \exp(\alpha_j + \beta^T \mathbf{x}_i)} - \frac{\exp(\alpha_{j-1} + \beta^T \mathbf{x}_i)}{1 + \exp(\alpha_{j-1} + \beta^T \mathbf{x}_i)} \right]^{y_{ij}} \right\}, \end{aligned} \quad (8.6)$$

viewed as a function of $(\{\alpha_j\}, \beta)$. This can be maximized to obtain the ML estimates using the Newton-Raphson algorithm (see McCullagh and Nelder, 1989) or the Newton-Raphson algorithm.

- The model parameters are estimated by maximum likelihood

Latent Variable Motivation for Proportional Odds Model

- A continuous latent variable y^* with $y^* = \tilde{\beta}^T \mathbf{x} + \epsilon$ and the distribution function of ϵ is $G(\cdot)$ (not mean zero)
- Thresholds $-\infty = \tilde{\alpha}_0 < \tilde{\alpha}_1 < \dots < \tilde{\alpha}_J = \infty$
- The observed response y satisfies $y = j$ if $\tilde{\alpha}_{j-1} < y^* \leq \tilde{\alpha}_j$


$$P(Y \leq j | \mathbf{x}) = P(y^* \leq \tilde{\alpha}_j | \mathbf{x}) = G(\tilde{\alpha}_j - \tilde{\beta}^T \mathbf{x})$$

- The proportional odds model

$$P(Y \leq j | \mathbf{x}) = \text{expit}(\alpha_j + \beta^T \mathbf{x})$$

- $G(\cdot) \sim \text{expit}(\cdot)$, $G^{-1}(\cdot) \sim \text{logit}(\cdot)$, $\tilde{\alpha}_j = \alpha_j$, $\tilde{\beta} = -\beta$

Proportional Odds Model: Interpretation using latent variable



Probit and Logit: Latent Variable Motivation for Binary Outcome Model

- A continuous latent variable y^* with $y^* = \tilde{\beta}^T \mathbf{x} + \epsilon$ (no intercept) and the distribution function of ϵ is $G(\cdot)$
- Threshold $-\infty < \tilde{\alpha} < \infty$. The observed response y satisfies $y = 1$ if $y^* \leq \tilde{\alpha}$ and $y = 0$ otherwise

$$P(Y = 1|\mathbf{x}) = P(y^* \leq \tilde{\alpha}|\mathbf{x}) = G(\tilde{\alpha} - \tilde{\beta}^T \mathbf{x})$$




- The logistic regression model: $P(Y = 1|\mathbf{x}) = \text{expit}(\alpha + \beta^T \mathbf{x})$.
 $G(\cdot) \sim \text{expit}(\cdot)$, $G^{-1}(\cdot) \sim \text{logit}(\cdot)$, $\tilde{\alpha} = \alpha$, $\tilde{\beta} = -\beta$
- The probit regression model: $P(Y = 1|\mathbf{x}) = \Phi(\alpha + \beta^T \mathbf{x})$.
 $G(\cdot) \sim \Phi(\cdot)$, $G^{-1}(\cdot) \sim \Phi^{-1}(\cdot)$, $\tilde{\alpha} = \alpha$, $\tilde{\beta} = -\beta$

Check the Proportional Odds Assumption

$$P(Y \leq j|\mathbf{x}) = \text{expit}(\alpha_j + \beta^T \mathbf{x})$$

- Proportional odds model is parsimonious and easy to interpret
- Replacing β with β_j may cause the cumulative probabilities to cross
- A score test of proportional odds model is available (SAS PROC LOGISTIC)
- Retain proportional odds model unless there is strong deviation from this assumption
- What to do when the proportional odds assumption is violated:
 - Adding additional terms, such as interaction
 - alternative ordinal model (next slides)
 - partial proportional odds model (SAS PROC LOGISTIC)
 - baseline category logit model

Alternative Models for Ordinal Data

- Cumulative link model: $G^{-1} [P(Y \leq j|\mathbf{x})] = \alpha_j + \beta^T \mathbf{x}$
- Cumulative probit model: $\Phi^{-1} [P(Y \leq j|\mathbf{x})] = \alpha_j + \beta^T \mathbf{x}$
- Cumulative complementary log-log model:
 $\log \{ -\log [1 - P(Y \leq j|\mathbf{x})] \} = \alpha_j + \beta^T \mathbf{x}$
 - Equivalent to a latent variable model with extreme value distribution for the residuals
 - Equivalent to proportional hazard model in for discrete survival data analysis (e.g., year of death at 1, 2, 3, ...) 
 - Rare event logistic regression
- Adjacent category logit model 
- Continuation ratio logit model 

Regression Models for Counts Data

Poisson Regression: introduction

- Applicable to counts data 0, 1, 2, Popularized in the 1970s and 1980s.
- Suppose Y has Poisson distribution with mean μ

$$P(Y = y) = \frac{e^{-\mu} \mu^y}{y!}, \quad y = 0, 1, 2, \dots \quad (1)$$

and $E(Y) = \text{var}(Y) = \mu$

- Suppose we want to relate Y to a covariate X , we assume $g(\mu) = \alpha + \beta X$. What choices do we have for the link function $g(\cdot)$? Any monotone mapping from $(0, \infty)$ to $(-\infty, \infty)$?
- Log link (log linear model) $\mu = \exp(\alpha + \beta X)$: the multiplicative effect of X on μ . Interpret $\exp(\beta)$ as a rate ratio.
- Example: A study of 400 patients with malignant melanoma. Y : tumor counts; X : tumor type (Hutchinson's, Superficial, Nodular, Indeterminate) and tumor site (Head & Neck, Trunk, Extremities)

Poisson Regression: Offset

- Frequently, discrete counts represent information collected over time (days, years) or in space (volume for bacteria counts) and interest lies in modeling rates. Denote the exposure time or volume by N , then the rate is Y/N , which has expectation μ/N .
- Modeling this rate with a log linear model as $\log \frac{\mu}{N} = \alpha + \beta X$. That is $\log \mu = \alpha + \beta X + \log(N)$ or $\mu = N \exp(\alpha + \beta X)$.
- $Y \sim \text{Poisson}(\mu)$.




- The log-likelihood is

$$L = \log \prod_{i=1}^n \mu_i^{Y_i} \{ \exp(-\mu_i) \} / Y_i! \quad , \quad \mu_i = N_i \exp(\alpha + \beta X_i)$$

Poisson Regression: Example with Offset

- Table 12.2 of “Categorical Data Analysis with the SAS System”.

Region	Age Group	Cases	Total
North	< 35	61	2,880,262
South	35 – 44	76	564,535
...



- Counts**: number of new cases in 1961-1971; **offset**: size of the populations at risk; **covariates**: age group and region
- Can this example be analyzed with binomial regression? 

```
proc genmod data = melanoma order = data ;  
class age region ;  
model cases = age region / dist = poisson link = log offset = ltotal ;  
run ;
```

Poisson Regression: Overdispersion

- Real data often have more variation than expected from a Poisson distribution (due to unobserved heterogeneity not captured by the covariates); $\text{var}(Y) > E(Y)$.
- Discover overdispersion through overdispersion parameter (next slide)
- Three ways to correct for overdispersion
 - Overdispersion model
 - Negative binomial regression
 - Poisson regression with random effect (later)


Overdispersion Model

- $\text{var}(Y) = \phi E(Y)$. $\phi > 1$ is the dispersion parameter. ($0 < \phi < 1$ is called under dispersion)
- $\mu_i = E(Y_i|\mathbf{X}_i) = N_i \exp(\mathbf{X}_i^T \theta)$ and $\text{var}(Y_i|\mathbf{X}_i) = \phi \mu_i$. In a Poisson regression model, we model the mean as a function of covariates, but not ϕ
- The likelihood equation $0 = \sum_{i=1}^n (Y_i - N_i \exp(\mathbf{X}_i^T \theta)) \mathbf{X}_i$, which uses only the mean function. So the point estimator is always correct even when overdispersion is ignored; but variance estimation and p-values will be incorrect.
- Estimate ϕ , and then multiply the ordinary standard error estimates by $\sqrt{\hat{\phi}}$. Example in page 150 
- An approximate method motivated from quasi-likelihood ideas
- Maybe wrong if the mean function is misspecified (major weakness)

Overdispersion with Binomial Model

- y_i is the sample proportion from n_i Bernoulli trials with parameter π_i , $i = 1, 2, \dots, n$.
- According to binomial model $E(Y_i) = \pi_i$ and $\text{var}(Y_i) = \pi_i(1 - \pi_i)/n_i$.
- $v(\pi_i) = \phi\pi_i(1 - \pi_i)/n_i$
- Estimate ϕ by $X^2/(n - p)$ and multiply the standard error by $\sqrt{(\hat{\phi})}$
- Not applicable when some $n_i = 1$ (and some not) because ϕ can only be 1 in this case.

Negative Binomial Regression §4.3.4

- If Y has a Poisson distribution with both mean and variance equal to λ , and λ has a gamma distribution (extra variation; overdispersion) with mean μ and variance μ^2/k , then marginally Y has a negative binomial distribution with mean μ and variance $\mu + \gamma\mu^2$ ($\gamma = 1/k > 0$) 
- When $\gamma \rightarrow 0$ (random effect variance approaches zero), the NB distribution approaches Poisson
- Like the overdispersion model, we let γ to be a overdispersion constant and model the mean $\mu = N \exp(\mathbf{X}^T \theta)$.
- Standard error will become larger after applying negative binomial regression
- Preferred over overdispersion model
- Available in SAS PROC GENMOD